

**Embargoed until: 17 July: 7am EST / 12pm UK / 1pm CET**



*PeerJ – the Journal of Life & Environmental Sciences*

## **PRESS RELEASE**

### **New Study Uncovers Taxonomic Breakthrough in the Common *Ophiuroid Ophiothrix angulata* (Echinodermata: Ophiuroidea)**



Invertebrate Zoology Collection, Florida Museum of Natural History, University of Florida, and Y. Quetzalli Hernández-Díaz.

*Ophiothrix angulata*, a widely recognized and prevalent ophiuroid species in the Western Atlantic, has long been the subject of taxonomic debate due to its remarkable morphological diversity. A new study just published in *PeerJ Life & Environment* has shed light on the species' taxonomy, revealing a significant scientific breakthrough.

Led by a team of researchers from Universidad Nacional Autónoma de México, Universidad Católica del Maule and Florida Natural History Museum, the comprehensive study aimed to assess species

delimitation and geographic differentiation within *O. angulata*. Quetzalli Hernández, co-author of the study, emphasizes the need for an integrative approach: "We tried to unravel the genetic relationship between the various shapes and colorations of the arm in the ophiuroid species *Ophiothrix angulata*. This species is widely distributed across different latitudes and depths, and its classification has caused confusion since its description almost 200 years ago, due to the great variety of colorations and subtle differences observed, which have confused alpha taxonomists who have relied solely on morphological characteristics. So, to address this taxonomic challenge, we adopted an integrative approach, conducting separate analyses on multiple types of genetic and morphological data before combining them. By compiling extensive evidence and adhering to species definition criteria, we have determined that *Ophiothrix angulata* represents a species complex with cryptic diversity more or less delimited within large geographic regions. Undertaking this task was undeniably challenging, yet equally exhilarating."

The study revealed significant discoveries regarding *Ophiothrix angulata*, which has traditionally been recognized as a single species. Hernández explained the findings, stating, "In the northwestern Atlantic distribution, we have identified two distinct genetic clades using the COI (mitochondrial) and ITS2 (nuclear) genetic markers. Notably, this genetic differentiation aligns with the variations observed in the shape of the ventral and dorsal arm plates. By combining genetic and morphometric data through integrative analyses, we have gathered substantial evidence to support the existence of more than one species within the cryptic complex of *Ophiothrix angulata*. One of these genetic clades reveals that a group previously classified as *O. angulata* belongs to a newly ophiuroid species. Additionally, an exhaustive analysis of the arm coloration patterns has provided scientific evidence indicating that coloration alone is not a reliable morphological characteristic for distinguishing the different genetic clades within our study. This finding holds great significance for the *O. angulata* species complex, as it challenges the validity of subspecies that were described solely based on coloration patterns in the 19th century, which remain relevant today."

The identification of multiple species within the *Ophiothrix angulata* complex demonstrates the importance of integrating various data types in taxonomic studies. This breakthrough not only enhances our understanding of the species' evolutionary history but also has implications for conservation efforts and management strategies.

The study serves as a milestone in the field of ophiuroid research and highlights the value of interdisciplinary approaches. It paves the way for further exploration into the ecology, behavior, and distribution patterns of these newly identified species within *Ophiothrix angulata*.



## Full Media Pack of article PDF, images

<https://drive.google.com/drive/folders/1yTmL59W68klOxyAZMjDUONHgstnCFsON?usp=sharing>

All media is CC BY 4.0. Photo/video credits are included in the filename.

**EMBARGOED - Embargoed until: 17 July: 7am EST / 12pm UK / 1pm CET**

**LINK TO THE PUBLISHED VERSION OF THIS ARTICLE:** <https://peerj.com/articles/15655/> The link will ONLY work after the embargo lifts. Your readers will be able to freely access this article via this URL.

### Cite this article

Hernández-Díaz YQ, Solis F, Beltrán-López RG, Benítez HA, Díaz-Jaimes P, Paulay G. 2023. Integrative species delimitation in the common ophiuroid *Ophiothrix angulata* (Echinodermata: Ophiuroidea): insights from COI, ITS2, arm coloration, and geometric morphometrics. *PeerJ* 11:e15655 <https://doi.org/10.7717/peerj.15655>

###

### About:

[PeerJ](#) is an Open Access publisher of seven peer-reviewed journals. PeerJ's mission is to give researchers the publishing tools and services they want with a unique and exciting experience. All works published by PeerJ are Open Access and published using a Creative Commons license (CC-BY 4.0). PeerJ is based in San Diego, CA and the UK and can be accessed at [peerj.com](https://peerj.com).

In 2022 PeerJ is celebrating its first decade of publishing and innovation.

*PeerJ* is the peer-reviewed journal for Biology, Medicine and Environmental Sciences. *PeerJ* has an Editorial Board of over 2,000 respected academics. PeerJ Media Resources (including logos) can be found at: [peerj.com/about/press](https://peerj.com/about/press)

###

### Media Contacts

For the authors:

Yoalli Quetzalli Hernández-Díaz [quetzalli.hernandez@gmail.com](mailto:quetzalli.hernandez@gmail.com)

For PeerJ:

Euan Lockie: [press@peerj.com](mailto:press@peerj.com)

*Note: If you would like to join the PeerJ Press Release list, please email your details to:*  
[press@peerj.com](mailto:press@peerj.com)